

**Hanoi University of Science and Technology**  
**School of Applied Mathematics and Informatics**  
**Exercises: Calculus 2**  
**Course ID: MI 1026**

## Chapter 1. Vectors and the geometry of space

**Exercise 1.** Determine whether the given vectors are orthogonal, parallel, or neither

(a)  $\vec{a} = (-5; 3; 7), \vec{b} = (6; -8; 2).$

(c)  $\vec{a} = -\vec{i} + 2\vec{j} + 5\vec{k}, \vec{b} = 3\vec{i} + 4\vec{j} - \vec{k}.$

(b)  $\vec{a} = (4; 6), \vec{b} = (-3; 2).$

(d)  $\vec{u} = (a, b, c), \vec{v} = (-b, a, 0).$

**Exercise 2.** For what values of  $b$  are the vectors  $(-6; b; 2)$  and  $(b; 2b; b)$  orthogonal?

**Exercise 3.** Find two unit vectors that make an angle of  $30^\circ$  with  $v = (3; 4).$

**Exercise 4.** Find the angle between a diagonal of a cube and one of its edges.

**Exercise 5.** Find an unit vector that parallel with  $8\vec{i} - \vec{j} + 4\vec{k}.$

**Exercise 6.** Find area of triangle  $ABC$ , where  $A(2; 8; 12), B(4; 5; 8), C(1; 4; 10).$

**Exercise 7.** Find the altitude  $AH$  of triangle  $ABC$ , where  $A(1; 6; 4), B(2; 5; 8)$  and  $C(-1; 4; 0).$

**Exercise 8.** Prove that  $\vec{x} \times (\vec{y} \times \vec{z}) = (\vec{x} \cdot \vec{z}) \cdot \vec{y} - (\vec{x} \cdot \vec{y}) \cdot \vec{z}.$

**Exercise 9.** Find the distance from the point  $(3; 7; -5)$  to

(a)  $xy$ -plane

(c)  $zx$ -plane

(e)  $y$ -axis

(b)  $yz$ -plane

(d)  $x$ -axis

(f)  $z$ -axis

**Exercise 10.** Evaluate  $\vec{a} + \vec{b}, 2\vec{a} + 3\vec{b}, |\vec{a}|$  and  $|\vec{a} - \vec{b}|.$

(a)  $\vec{a} = 4\vec{i} + \vec{j}$  and  $\vec{b} = \vec{i} - 2\vec{j}.$

(b)  $\vec{a} = \vec{i} + 2\vec{j} - 3\vec{k}$  and  $\vec{b} = -2\vec{i} - \vec{j} + 5\vec{k}.$

**Exercise 11.** Find the angle between

(a)  $\vec{a} = (3; 4)$  and  $\vec{b} = (5; 12).$

(b)  $\vec{a} = (6; -3; 2)$  and  $\vec{b} = (2; 1; -2).$

**Exercise 12.** Find a unit vector that is orthogonal to both  $\vec{i} + \vec{j}$  and  $\vec{i} + \vec{k}.$

**Exercise 13.** Find the angle between a diagonal of a cube and a diagonal of one of its faces.

**Exercise 14.** Prove that  $|\vec{a} + \vec{b}|^2 + |\vec{a} - \vec{b}|^2 = 2|\vec{a}|^2 + 2|\vec{b}|^2.$

**Exercise 15.** Find the volume of the parallelepiped determined by the vectors

(a)  $\vec{a} = (6; -3; -1)$ ,  $\vec{b} = (0; 1; 2)$  and  $\vec{c} = (4; -2; 5)$ .

(b)  $\vec{a} = \vec{i} + \vec{j} - \vec{k}$ ,  $\vec{b} = \vec{i} - \vec{j} + \vec{k}$  and  $\vec{c} = -\vec{i} + \vec{j} + \vec{k}$ .

**Exercise 16.** (a) Let  $P$  be a point not on the line that passes through the points  $Q, R$  and  $S$ . Show that the distance  $d$  from the point  $P$  to the plane  $(QRS)$  is

$$d = \frac{|\vec{a} \cdot (\vec{b} \times \vec{c})|}{|\vec{a} \times \vec{b}|},$$

where  $\vec{a} = \overrightarrow{QR}$ ,  $\vec{b} = \overrightarrow{QS}$  and  $\vec{c} = \overrightarrow{QP}$ .

(b) Use the formula in part (a) to find the distance from the point  $P(2; 1; 4)$  to the plane through the points  $Q(1; 0; 0)$ ,  $R(0; 2; 0)$  and  $S(0; 0; 3)$ .

**Exercise 17.** Find an equation of the sphere with center  $(1; -4; 3)$  and radius 5. Describe its intersection with each of the coordinate planes.

**Exercise 18.** Find an equation of the sphere that passes through the origin and whose center is  $(1; 2; 3)$ .

**Exercise 19.** Find an equation of a sphere if one of its diameters has end points  $(2; 1; 4)$  and  $(4; 3; 10)$ .

**Exercise 20.** Find an equation of the largest sphere with center  $(5; 4; 9)$  that is contained in the first octant.

**Exercise 21.** Consider the points  $P$  such that the distance from  $P$  to  $A(-1; 5; 3)$  is twice the distance from  $P$  to  $B(6; 2; -2)$ . Show that the set of all such points is a sphere, and find its center and radius.

**Exercise 22.** Find an equation of the set of all points equidistant from the points  $A(-1; 5; 3)$  and  $B(6; 2; -2)$ . Describe the set.

**Exercise 23.** Sketch and classify the quadric surface

(a)  $x^2 + 2z^2 - 6x - y + 10 = 0$ .      (c)  $x^2 = y^2 + 4z^2$ .      (e)  $y = z^2 - x^2$ .

(b)  $x = y^2 + 4z^2$ .      (d)  $-x^2 + 4y^2 - z^2 = 4$ .      (f)  $4x^2 - 16y^2 + z^2 = 16$ .

**Exercise 24.** Find an equation for the surface obtained by rotating

(a) the parabol  $y = x^2$  about the  $y$ -axis.      (b) the line  $x = 3y$  about the  $x$ -axis.

**Exercise 25.** Find an equation for the surface consisting of all points that are equidistant from the point  $(-1; 0; 0)$  and the plane  $x = 1$ . Identify the surface.

**Exercise 26.** Find an equation for the surface consisting of all points  $P$  for which the distance from  $P$  to the  $x$ -axis is twice the distance from  $P$  to the  $yz$ -plane. Identify the surface.

**Exercise 27.** Change from rectangular to cylindrical coordinates.

(a)  $(-1; 1; 1)$ .

(c)  $(2\sqrt{3}; 2; -1)$ .

(b)  $(-2; 2\sqrt{3}; 3)$ .

(d)  $(4; -3; 2)$ .

**Exercise 28.** Write the equations in cylindrical coordinates.

(a)  $x^2 - x + y^2 + z^2 = 1$ .

(c)  $3x + 2y + z = 6$ .

(b)  $z = x^2 - y^2$ .

(d)  $-x^2 - y^2 + z^2 = 1$ .

**Exercise 29.** Identify the surface whose equation is given

(a)  $z = 4 - r^2$ .

(b)  $2r^2 + z^2 = 1$ .

**Exercise 30.** Change from rectangular to spherical coordinates

(a)  $(0; -2; 0)$ .

(c)  $(1; 0; \sqrt{3})$ .

(b)  $(-1; 1; -\sqrt{2})$ .

(d)  $(\sqrt{3}; -1; 2\sqrt{3})$ .

**Exercise 31.** Identify the surface whose equation is given.

(a)  $r = \sin \theta \sin \varphi$ .

(b)  $r^2(\sin^2 \varphi \sin^2 \theta + \cos^2 \varphi) = 9$ .

**Exercise 32.** Write the equation in spherical coordinates

(a)  $z^2 = x^2 + y^2$ .

(c)  $x^2 - 2x + y^2 + z^2 = 0$ .

(b)  $x^2 + z^2 = 9$ .

(d)  $x + 2y + 3z = 1$ .

## Chapter 2. Vector Functions

**Exercise 33.** Find the parametric equations for the intersection of the circular cylinder  $x^2 + y^2 = 4$  and parabolic cylinder  $z = x^3$ .

**Exercise 34.** Find the domain.

(a)  $\vec{r}(t) = (\sqrt{4 - t^2}, e^{-3t}, \ln(1 + t))$ .

(c)  $\vec{r}(t) = \arcsin \frac{2t}{1+t} \vec{i} + \frac{\sqrt{t}}{\sin \pi t} \vec{k}$ .

(b)  $\vec{r}(t) = \frac{t-2}{t+2} \vec{i} + \sin t \vec{j} + \ln(9 - t^2) \vec{k}$ .

(d)  $\vec{r}(t) = (\sqrt{\cosh t - 1}, \sqrt{t^4 - 5t^2 + 4}, 0)$ .

**Exercise 35.** Find the limit

$$(a) \lim_{t \rightarrow 0} \left( \frac{e^t - 1}{t}, \frac{\sqrt{t+1} - 1}{t}, \frac{3}{t+1} \right). \quad (b) \lim_{t \rightarrow +\infty} \left( \arctan t, e^{-2t}, \frac{\ln t}{t+1} \right).$$

**Exercise 36.** Find a vector function that represents the curve of intersection of the two surfaces.

(a) The cylinder  $x^2 + y^2 = 4$  and the surface  $z = xy$ .

(b) The paraboloid  $z = 4x^2 + y^2$  and the parabolic cylinder  $y = x^2$ .

**Exercise 37.** Suppose  $u$  and  $v$  are vector functions that possess limits as  $t \rightarrow a$  and let  $c$  be a constant. Prove the following properties of limits.

$$(a) \lim_{t \rightarrow a} [\vec{u}(t) + \vec{v}(t)] = \lim_{t \rightarrow a} \vec{u}(t) + \lim_{t \rightarrow a} \vec{v}(t). \quad (c) \lim_{t \rightarrow a} [\vec{u}(t) \cdot \vec{v}(t)] = \lim_{t \rightarrow a} \vec{u}(t) \cdot \lim_{t \rightarrow a} \vec{v}(t).$$

$$(b) \lim_{t \rightarrow a} c\vec{v}(t) = c \lim_{t \rightarrow a} \vec{v}(t). \quad (d) \lim_{t \rightarrow a} [\vec{u}(t) \times \vec{v}(t)] = \lim_{t \rightarrow a} \vec{u}(t) \times \lim_{t \rightarrow a} \vec{v}(t).$$

**Exercise 38.** Find the derivative of the vector function.

$$(a) \vec{r}(t) = (t \sin t, t^3, t \cos 2t).$$

$$(c) \vec{r}(t) = e^{t^2} \vec{i} - \sin^2 t \vec{j} + \ln(1 + 3t) \vec{k}.$$

$$(b) \vec{r}(t) = \arcsin t \vec{i} + \sqrt{1 - t^2} \vec{j} + \vec{k}.$$

$$(d) \vec{r}(t) = (e^{\sin t}, \arctan t, t^2).$$

**Exercise 39.** Find parametric equations for the tangent line to the curve with the given parametric equations at the specified point.

$$(a) \begin{cases} x = t, \\ y = e^{-t}, \\ z = 2t - t^2, \end{cases} \quad \text{at } (0; 1; 0).$$

$$(b) \begin{cases} x = t \cos t, \\ y = t, \\ z = t \sin t, \end{cases} \quad \text{at } (-\pi; \pi; 0).$$

**Exercise 40.** Find the point of intersection of the tangent lines to the curve  $\vec{r}(t) = (\sin \pi t; 2 \sin \pi t; \cos \pi t)$  at the points where  $t = 0$  and  $t = 0.5$ .

**Exercise 41.** Evaluate the integral.

$$(a) \int_0^{\pi/2} (3 \sin^2 t \cos t \vec{i} + 3 \sin t \cos^2 t \vec{j} + 2 \sin t \cos t \vec{k}) dt.$$

$$(b) \int_1^2 (t^2 \vec{i} + t\sqrt{t-1} \vec{j} + t \sin \pi t \vec{k}) dt.$$

$$(c) \int_1^2 (e^t \vec{i} + 2t \vec{j} + \ln t \vec{k}) dt.$$

$$(d) \int_0^{1/4} (\cos \pi t \vec{i} + \sin \pi t \vec{j} + t^2 \vec{k}) dt.$$

**Exercise 42.** If a curve has the property that the position vector  $\vec{r}(t)$  is always perpendicular to the tangent vector  $\frac{d}{dt} \vec{r}(t)$ , show that the curve lies on a sphere with center the origin.

**Exercise 43.** Find the length of the curve.

(a)  $\vec{r}(t) = (2 \sin t, 5t, 2 \cos t); -10 \leq t \leq 10.$

(c)  $\vec{r}(t) = (\cos t, \sin t, \ln \cos t); 0 \leq t \leq \pi/4.$

(b)  $\vec{r}(t) = (2t, t^2, \frac{1}{3}t^3); 0 \leq t \leq 1.$

(d)  $\vec{r}(t) = (\sin t - t \cos t, \cos t + t \sin t, t^2); 0 \leq t \leq 2\pi.$

**Exercise 44.** Let  $C$  be the curve of intersection of the parabolic cylinder  $x^2 = 2y$  and the surface  $3z = xy$ . Find the length of  $C$  from the origin to the point  $(6; 18; 36)$ .

**Exercise 45.** Suppose you start at the point  $(0; 0; 3)$  and move 5 units along the curve  $x = 3 \sin t; y = 4t; z = 3 \cos t$  in the positive direction. Where are you now?

**Exercise 46.** Find the curvature of  $\vec{r}(t) = (e^t \cos t, e^t \sin t, t)$  at the point  $(1; 0; 0)$ .

**Exercise 47.** Find the curvature of  $\vec{r}(t) = (t, t^2, t^3)$  at the point  $(1; 1; 1)$ .

**Exercise 48.** Find the curvature of  $y = \sqrt{x^2 + 1} - 2$  at the point  $A(0; -1)$ .

**Exercise 49.** Find the curvature of the curve given by  $\begin{cases} x^2 + y^2 + 1 = 2(x - y), \\ x + y - z = 2 \end{cases}$  at the point  $A(1; 0; -1)$ .

**Exercise 50.** Find the curvature.

(a)  $\vec{r}(t) = t\vec{i} + t\vec{j} + (1 + t^2)\vec{k}.$

(c)  $x = e^t \cos t, y = e^t \sin t.$

(b)  $\vec{r}(t) = 3t\vec{i} + 4 \sin t\vec{j} + 4 \cos t\vec{k}.$

(d)  $x = t^3 + 1, y = t^2 + 1.$

**Exercise 51.** Find the curvature.

(a)  $y = 2x - x^2.$

(c)  $y = 4x^{5/2}.$

(b)  $y = \cos x.$

(d)  $y = \sin x.$

**Exercise 52.** At what point does the curve have maximum curvature? What happens to the curvature as  $x \rightarrow \infty$ .

(a)  $y = \ln x.$

(b)  $y = e^x.$

**Exercise 53.** Find an equation of a parabola that has curvature 4 at the origin.

**Exercise 54.** Find the velocity vector, acceleration vector, and speed of a moving particle with the given position function

(a)  $\vec{r}(t) = (e^{-t}, t\sqrt{3}, e^t).$

(b)  $\vec{r}(t) = e^t(\sin t, t, \cos t).$

**Exercise 55.** A moving particle starts at an initial position  $\vec{r}(0) = (2; -3; 4)$  with initial velocity  $\vec{v}(0) = (1; 5; -4)$ . Find its velocity and position at time  $t$  if  $\vec{a}(t) = (2, t^3, e^{3t})$ . Find its speed at  $t = 1$ .

## Chapter 3. Double Integrals

**Exercise 56.** Find the volume of the solid that lies under the plane  $4x + 6y - 2z + 15 = 0$  and above the rectangle  $R = \{(x, y) : -1 \leq x \leq 2, -1 \leq y \leq 1\}$ .

**Exercise 57.** Find the volume of the solid enclosed by the surface  $z = 1 + e^x \sin y$  and the planes  $x = \pm 1$ ,  $y = 0$ ,  $y = \pi$  and  $z = 0$ .

**Exercise 58.** Find the volume of the solid in the first octant bounded by the cylinder  $z = 16 - x^2$  and the plane  $y = 5$ .

**Exercise 59.** Find the volume of the solid enclosed by the paraboloid  $z = 2 + x^2 + (y - 2)^2$  and the planes  $z = 1$ ,  $x = 1$ ,  $x = -1$ ,  $y = 0$ , and  $y = 4$ .

**Exercise 60.** Evaluate the following integrals

(a)  $\int_1^3 dx \int_1^5 \frac{\ln y}{xy} dy.$

(b)  $\int_0^1 dx \int_0^1 xy \sqrt{x^2 + y^2} dy.$

(c)  $\iint_R \frac{1}{1+x+y} dx dy$ , where  $R = [1; 3] \times [1; 2]$ .

(d)  $\iint_D ye^{xy} dx dy$ , where  $D = \{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq 2; 0 \leq y \leq 3\}$ .

(e)  $\iint_D |x+y| dx dy$ , where  $D = \{(x, y) \in \mathbb{R}^2 : |x| \leq 1; |y| \leq 1\}$ .

(f)  $\iint_D \sqrt{|y-x^2|} dx dy$ , where  $D = \{(x, y) \in \mathbb{R}^2 : |x| \leq 1, 0 \leq y \leq 2\}$ .

(g)  $\iint_D |y-x^2|^3 dx dy$ , where  $D = \{(x, y) \in \mathbb{R}^2 : |x| \leq 1, \text{ and } 0 \leq y \leq 2\}$ .

**Exercise 61.** Use Midpoint rule to estimate the volume of the solid that lies below the surface  $z = xy$  and above the rectangle  $R = \{(x, y) : 0 \leq x \leq 6, 0 \leq y \leq 4\}$ . Use a Riemann sum with  $m = 3, n = 2$ .

**Exercise 62.**

(a) Estimate the volume of the solid that lies below the surface  $z = 1 + x^2 + 3y$  and above the rectangle  $R = [1; 2] \times [0; 3]$ . Use a Riemann sum with  $m = n = 2$  and choose the sample points to be lower left corners.

(b) Use the Midpoint Rule to estimate the volume in part (1).

**Exercise 63.** A table of values is given for a function  $f(x, y)$  defined on  $R = [0; 4] \times [2; 4]$ .

(a) Estimate  $\iint_R f(x, y) dx dy$  using Midpoint rule with  $m = n = 2$ .

(b) Estimate the double integral with  $m = n = 4$  by choosing the sample points to be the points closest to the origin.

$x \backslash y$	2.0	2.5	3.0	3.5	4.0
0	-3	-5	-6	-4	-1
1	-1	-2	-3	-1	1
2	1	0	-1	1	4
3	2	2	1	3	7
4	3	4	2	5	9

**Exercise 64.** Change the order of the following integration.

$$(a) \int_{-1}^1 dx \int_{-\sqrt{1-x^2}}^{1-x^2} f(x, y) dy.$$

$$(d) \int_0^{\frac{\pi}{2}} dy \int_{\sin y}^{1+y^2} f(x, y) dx.$$

$$(b) \int_0^1 dy \int_{2-y}^{1+\sqrt{1-y^2}} f(x, y) dx.$$

$$(e) \int_0^{\sqrt{2}} dy \int_0^y f(x, y) dx + \int_{\sqrt{2}}^2 dy \int_0^{\sqrt{4-y^2}} f(x, y) dx.$$

$$(c) \int_0^2 dx \int_{\sqrt{2x-x^2}}^{\sqrt{2x}} f(x, y) dy.$$

**Exercise 65.** Evaluate the integrals

$$(a) \int_0^1 dx \int_0^{1-x^2} \frac{xe^{3y}}{1-y} dy.$$

$$(b) \iint_D x^2(y-x) dx dy, \text{ where } D \text{ is bounded by } y = x^2 \text{ and } x = y^2.$$

$$(c) \iint_D \frac{y}{1+x^2} dx dy, \text{ where } D \text{ is bounded by } y = \sqrt{x}, y = 0 \text{ and } x = 1.$$

$$(d) \iint_D xy dx dy, \text{ where } D \text{ is bounded by } x = y^2, x = -1, y = 0 \text{ and } y = 1.$$

$$(e) \iint_D (x+y) dx dy, \text{ where } D \text{ is bounded by } x^2 + y^2 \leq 1, \sqrt{x} + \sqrt{y} \geq 1.$$

$$(f) \iint_D (x^2 + y^2)^{3/2} dx dy, \text{ where } D \text{ is a region in the first quadrant and bounded by } y = 0, y = \sqrt{3}x \text{ and circle } x^2 + y^2 = 9.$$

$$(g) \iint_D (|x| + |y|) dx dy, D = \{(x, y) \in \mathbb{R}^2 : |x| + |y| \leq 1\}.$$

**Exercise 66.** Change to polar coordinates in a double integral  $\iint_D f(x, y) dx dy$ , where  $D$  is a region as follows:

$$(a) a^2 \leq x^2 + y^2 \leq b^2.$$

$$(c) x^2 + y^2 \leq 2x, x^2 + y^2 \leq 2y.$$

$$(b) x^2 + y^2 \geq 4x, x^2 + y^2 \leq 8x, y \geq x, y \leq \sqrt{3}x.$$

**Exercise 67.** Use polar coordinates to find the following integrals

- (a)  $\int_0^R dx \int_0^{\sqrt{R^2-x^2}} \ln(1+x^2+y^2)dy, \quad (R > 0).$
- (b)  $\iint_D xy dx dy$ , where  $D$  is a disk  $(x-2)^2 + y^2 \leq 1, y \geq 0$ .
- (c)  $\iint_D (\sin y + 3x) dx dy$ , where  $D$  is a disk  $(x-2)^2 + y^2 \leq 1$ .
- (d)  $\iint_D |x+y| dx dy$ , where  $D$  is a disk  $x^2 + y^2 \leq 1$ .

**Exercise 68.** Evaluate the following integrals:

- (a)  $\iint_D \frac{2xy+1}{\sqrt{1+x^2+y^2}} dx dy$ , with  $D: x^2 + y^2 \leq 1$ .
- (b)  $\iint_D \frac{dx dy}{(x^2+y^2)^2}$ , with  $D: \begin{cases} y \leq x^2 + y^2 \leq 2y \\ x \leq y \leq \sqrt{3}x. \end{cases}$

**Exercise 69.** Find the mass and center of mass of the lamina that occupies the region  $D$  and has the given density function  $f(x, y)$ .

- (a)  $D = \{(x, y) : 1 \leq x \leq 3, 1 \leq y \leq 4\}, f(x, y) = 2y^2$ .
- (b)  $D = \{(x, y) : 0 \leq x \leq a, 0 \leq y \leq b\}, f(x, y) = 1 + x^2 + y^2$ .
- (c)  $D$  is bounded by  $y = 1 - x^2$  and  $y = 0$ ;  $f(x, y) = ky$ .
- (d)  $D$  is bounded by  $y = x^2$  and  $y = x + 2$ ;  $f(x, y) = kx$ .
- (e)  $D = \{(x, y) : 0 \leq y \leq \sin \frac{\pi x}{L}, 0 \leq x \leq L\}; f(x, y) = y$ .
- (f)  $D$  is bounded by the parabolas  $y = x^2$ , and  $x = y^2$ ;  $f(x, y) = \sqrt{x}$ .

**Exercise 70.** Find the area of the surface.

- (a) The part of the plane  $z = 2 + 3x + 4y$  that lies above the rectangle  $[0; 5] \times [1; 4]$ .
- (b) The part of the plane  $2x + 5y + z = 10$  that lies inside the cylinder  $x^2 + y^2 = 9$ .
- (c) The part of the paraboloid  $z = 4 - x^2 - y^2$  that lies above  $xy$ -plane.
- (d) The part of the sphere  $x^2 + y^2 + z^2 = 4z$  that lies inside the paraboloid  $z = x^2 + y^2$ .

**Exercise 71.** Use the change of variables  $u = x + y$  and  $v = x - y$  to evaluate the integral

$$\int_0^1 dx \int_{-x}^x (2-x-y)^2 dy.$$

**Exercise 72.** Evaluate the following integrals:

- (a)  $\iint_D \frac{xy}{x^2+y^2} dx dy$ , where  $D: \begin{cases} 2x \leq x^2 + y^2 \leq 12 \\ x^2 + y^2 \geq 2\sqrt{3}y \\ x \geq 0, y \geq 0. \end{cases}$



(b)  $\iint_D |9x^2 - 4y^2| dx dy$ , where  $D : \frac{x^2}{4} + \frac{y^2}{9} \leq 1$ .

## Chapter 4. Triple Integrals

**Exercise 73.** Express the triple integral  $\int_0^1 \int_{\sqrt{x}}^1 \int_0^{1-y} f(x, y, z) dz dy dx$  in the order  $dx dy dz$ .

**Exercise 74.** Evaluate the iterated integral

(a)  $\int_0^1 \int_x^{2x} \int_0^y 2xyz dz dy dx$ ,

(c)  $\int_0^1 \int_0^z \int_0^y ze^{-y^2} dx dy dz$ ,

(b)  $\int_0^3 \int_0^1 \int_0^{\sqrt{1-z^2}} ze^y dx dz dy$ ,

(d)  $\int_0^{\frac{\pi}{2}} \int_0^y \int_0^x \cos(x + y + z) dz dx dy$ .

**Exercise 75.** Evaluate the triple integral

(a)  $\iiint_E y dx dy dz$ , where  $E$  is bounded by the planes  $x = 0, y = 0, z = 0$  and  $2x + 2y + z = 4$ .

(b)  $\iiint_E x^2 e^y dx dy dz$ , where  $E$  is bounded by the parabolic cylinder  $z = 1 - y^2$  and the planes  $z = 0, x = 1, x = -1$ .

(c)  $\iiint_E xy dx dy dz$ , where  $E$  is bounded by the parabolic cylinder  $y = x^2, x = y^2$  and the planes  $z = 0, z = x + y$ .

(d)  $\iiint_E x dx dy dz$ , where  $E$  is bounded by the paraboloid  $x = 4y^2 + 4z^2$  and the plane  $x = 4$ .

(e)  $\iiint_E (x^3 + xy^2) dx dy dz$ , where  $E$  is the solid in the first octant that lies beneath the paraboloid  $z = 1 - x^2 - y^2$ .

**Exercise 76.** Find the volume of the region  $E$  bounded by the paraboloids  $z = x^2 + y^2$  and  $z = 36 - 3x^2 - 3y^2$ .

**Exercise 77.** Find the volume of the solid that lies within both the cylinder  $x^2 + y^2 = 1$  and the sphere  $x^2 + y^2 + z^2 = 4$ .

**Exercise 78.** Find the center of mass and the moments of inertia of the cubic  $[1; 2] \times [1; 2] \times [1; 2]$  if the density is  $\rho(x, y, z) = x^2 + y^2 + z^2$ .

**Exercise 79.** Find the center of mass and the moments of inertia of the tetrahedron with vertices  $(0; 0; 0); (1; 0; 0); (0; 1; 0)$  and  $(0; 0; 1)$  if the density is  $C$  (constant  $\neq 0$ ).

**Exercise 80.** Evaluate the integrals by changing to cylindrical coordinates.

$$(a) \int_{-2}^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \int_{\sqrt{x^2+y^2}}^2 xzdzdxdy,$$

$$(b) \int_{-3}^3 \int_0^{\sqrt{9-x^2}} \int_0^{9-x^2-y^2} \sqrt{x^2+y^2} dzdydx.$$

**Exercise 81.** Evaluate the triple integrals.

$$(a) \iiint_B x dx dy dz, \text{ where } B \text{ is bounded by the cone } x = \sqrt{y^2 + z^2}, \text{ and the plane } x = 1.$$

$$(b) \iiint_B \sqrt{x^2 + 4z^2} dx dy dz, \text{ where } B \text{ is bounded by the cone } x^2 + 4z^2 = y^2, \text{ and the plane } y = -1.$$

**Exercise 82.** Evaluate the triple integrals.

$$(a) \iiint_A \sqrt{x^2 + 4y^2 + z^2} dx dy dz, \text{ where } A \text{ is given by } x^2 + 4y^2 + z^2 \leq 2x.$$

$$(b) \iiint_A x^2 dx dy dz, \text{ where } A \text{ is bounded by the } xz\text{-plane and the hemispheres } y = \sqrt{9 - x^2 - z^2} \text{ and } y = \sqrt{16 - x^2 - z^2}.$$

**Exercise 83.** Find the moments of inertia of the ball  $B = \{x^2 + y^2 + z^2 \leq 1\}$  if the density  $\rho(x, y, z) = \sqrt{x^2 + y^2 + z^2}$ .

**Exercise 84.** Evaluate the integral by changing to spherical coordinates  $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_{x^2+y^2}^{\sqrt{2-x^2-y^2}} xydzdydx$ .

**Exercise 85.** Let  $E$  be the solid given by  $x \leq x^2 + y^2 + z^2 \leq 2x$ ,  $y \leq x^2 + y^2 + z^2 \leq 2y$ , and  $z \leq x^2 + y^2 + z^2 \leq 2z$ .

(a) Evaluate the Jacobian of doing change the variables:

$$u = \frac{x}{x^2 + y^2 + z^2}, \quad v = \frac{y}{x^2 + y^2 + z^2}, \quad w = \frac{z}{x^2 + y^2 + z^2}.$$

(b) Evaluate the triple integral  $\iiint_E \frac{1}{(x^2 + y^2 + z^2)^2} dx dy dz$ .

**Exercise 86.** Let  $E$  be the solid given by  $|x - y| + |x + 3y| + |x + y + z| \leq 1$ . Evaluate the triple integral  $\iiint_E xy dx dy dz$ .

## Chapter 5. Line Integrals

**Exercise 87.** Evaluate the following line integrals:

$$(a) \int_C xy ds, \text{ where } C : x = t^2, y = 2t, 0 \leq t \leq 1.$$

$$(b) \int_C xy^4 ds, \text{ where } C : x^2 + y^2 = 9, x \geq 0.$$

- (c)  $\int_C (x^2y^3 - \sqrt{x})dy$ , where  $C$  is the arc of the curve  $y = \sqrt{x}$  from  $(1; 1)$  to  $(4; 2)$ .
- (d)  $\int_C x^2dx + y^2dy$ , where  $C$  consists of circle  $x^2 + y^2 = 4$  from  $(2; 0)$  to  $(0; 2)$  and the segment from  $(0; 2)$  to  $(4; 3)$ .
- (e)  $\int_C (3x - y)ds$ , where  $C$  is the half of circle  $y = \sqrt{9 - x^2}$ .
- (f)  $\int_C (x - y)ds$ , where  $C$  is a circle  $x^2 + y^2 = 2x$ .
- (g)  $\int_C y^2ds$ , where  $C$  is given by  $x = a(t - \sin t)$ ,  $y = a(1 - \cos t)$  with  $0 \leq t \leq 2\pi$ ,  $a > 0$ .
- (h)  $\int_C \sqrt{x^2 + y^2}ds$ , where  $C$  is a curve  $x = a(\cos t + t \sin t)$ ,  $y = a(\sin t - t \cos t)$ , with  $0 \leq t \leq 2\pi$ ,  $a > 0$ .

**Exercise 88.** Evaluate the following line integrals:

- (a)  $\int_C (x^2 + y^2 + z^2)ds$ , where  $C : x = t, y = \cos 2t, z = \sin 2t, 0 \leq t \leq 2\pi$ .
- (b)  $\int_C xe^{yz}ds$ , where  $C$  is the segment from  $(0; 0; 0)$  to  $(1; 2; 3)$ .
- (c)  $\int_C ydx + zdy + xdz$ , where  $C : x = \sqrt{t}, y = t, z = t^2, 1 \leq t \leq 4$ .
- (d)  $\int_C (y + z)dx + (x + z)dy + (x + y)dz$ , where  $C$  consists of two line segments from  $(0; 0; 0)$  to  $(1; 0; 1)$ , and from  $(1; 0; 1)$  to  $(0; 1; 2)$ .

**Exercise 89.** Evaluate the following line integrals

- (a)  $\int_{AB} (x^2 - 2xy)dx + (2xy - y^2)dy$ , where  $AB$  is a part of parabol  $y = x^2$  from  $A(1; 1)$  to  $B(2; 4)$ .
- (b)  $\int_C (2x - y)dx + xdy$ , where  $C$  is a curve  $\begin{cases} x = a(t - \sin t) \\ y = a(1 - \cos t) \end{cases}$  whose direction is increasing direction of the parameter  $t$ , ( $0 \leq t \leq 2\pi, a > 0$ ).
- (c)  $\int_{ABCA} 2(x^2 + y^2)dx + x(4y + 3)dy$ , where  $ABCA$  is a broken line through the points  $A(0; 0)$ ,  $B(1; 1)$ ,  $C(0; 2)$ .
- (d)  $\int_{ABCD} \frac{dx + dy}{|x| + |y|}$ , where  $ABCD$  is a broken line through the points  $A(1; 0)$ ,  $B(0; 1)$ ,  $C(-1; 0)$ ,  $D(0; -1)$ .
- (e)  $\int_C \frac{\sqrt[4]{x^2 + y^2}dx}{2} + dy$ , where  $C$  is curve  $\begin{cases} x = t \sin \sqrt{t} \\ y = t \cos \sqrt{t}, (0 \leq t \leq \frac{\pi^2}{4}) \end{cases}$ .

**Exercise 90.** Evaluate the following line integral

$$\int_C (xy + x + y)dx + (xy + x - y)dy$$

in two ways: by computing it directly, and by Green's formula, then compare the results, where  $C$  is a curve:

(a)  $x^2 + y^2 = 2x$

(b)  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, (a, b > 0)$

**Exercise 91.** Evaluate the following line integrals:

(a)  $\oint_{x^2+y^2=2x} x^2 \left(y + \frac{x}{4}\right) dy - y^2 \left(x + \frac{y}{4}\right) dx.$

(b)  $\int_{OABO} e^x [(1 - \cos y)dx - (y - \sin y)dy],$  where  $OABO$  is a broken line through the points  $O(0; 0), A(1; 1), B(0; 2).$

(c)  $\oint_{x^2+y^2=2x} (xy + e^x \sin x + x + y)dx - (xy - e^{-y} + x - \sin y)dy.$

(d)  $\int_C (xy^4 + x^2 + y \cos(xy))dx + \left(\frac{x^3}{3} + xy^2 - x + x \cos(xy)\right) dy,$  where  $C$  is a curve  $x = a \cos t, y = a \sin t,$  ( $a > 0$ ).

(e)  $\oint_C (e^x + y^6)dx + (e^y + 3x)dy,$  where  $C$  is a boundary of region enclosed by  $x = 14 + \sqrt{|y|}$  and  $x = y^2,$  with oriented counterclockwise.

**Exercise 92.** Using the line integral of the second kind in order to compute the area of the region bounded by an arch of the cycloid:  $x = a(t - \sin t), y = a(1 - \cos t)$  and  $x$ -axis, ( $a > 0$ ).**Exercise 93.** Evaluate the following line integral

(a)  $\int_{(-2;-1)}^{(3;0)} (x^4 + 4xy^3)dx + (6x^2y^2 - 5y^4)dy.$

(b)  $\int_{(1;\pi)}^{(2;2\pi)} \left(1 - \frac{y^2}{x^2} \cos \frac{y}{x}\right)dx + \left(\sin \frac{y}{x} + \frac{y}{x} \cos \frac{y}{x}\right)dy.$

**Exercise 94.** Evaluate the line integral

$$I = \int_L \left(3x^2y^2 + \frac{2}{4x^2 + 1}\right)dx + \left(3x^3y + \frac{2}{y^3 + 4}\right)dy,$$

where  $L$  is curve  $y = \sqrt{1 - x^4}$  from  $A(1; 0)$  to  $B(-1; 0).$ **Exercise 95.** Find the constant  $\alpha$  such that the following integral is an independent of path in the domain

$$\int_{AB} \frac{(1 - y^2)dx + (1 - x^2)dy}{(1 + xy)^\alpha}.$$

**Exercise 96.** Find the curl and the divergence of the vector

(a)  $\vec{F}(x, y, z) = xy\vec{i} + yz\vec{j} + zx\vec{k}.$

(b)  $\vec{F}(x, y, z) = \frac{x}{x^2 + y^2 + z^2}\vec{i} + \frac{y}{x^2 + y^2 + z^2}\vec{j} + \frac{z}{x^2 + y^2 + z^2}\vec{k}.$

**Exercise 97.** Prove that

$$(a) \operatorname{curl}(\vec{F} + \vec{G}) = \operatorname{curl}\vec{F} + \operatorname{curl}\vec{G}.$$

$$(b) \operatorname{curl}(f\vec{F}) = f\operatorname{curl}\vec{F} + (\nabla f) \times \vec{F}.$$

**Exercise 98.** Determine whether or not  $\vec{F}$  is a conservative vector field. If it is, find a function  $f$  such that  $\vec{F} = \nabla f$ .

$$(a) \vec{F}(x, y) = (2x - 3y)\vec{i} + (-3x + 4y - 8)\vec{j}.$$

$$(b) \vec{F}(x, y) = e^x \cos y\vec{i} + e^x \sin y\vec{j}.$$

$$(c) \vec{F}(x, y) = (xy \cos(xy) + \sin(xy))\vec{i} + (x^2 \cos(xy))\vec{j}.$$

$$(d) \vec{F}(x, y) = (\ln y + 2xy^3)\vec{i} + (3x^2y^2 + \frac{x}{y})\vec{j}.$$

**Exercise 99.** Find  $f$  such that  $\vec{F} = \nabla f$  and then compute  $\int_C \vec{F} \cdot d\vec{r}$ .

$$(a) \vec{F}(x, y) = xy^2\vec{i} + x^2y\vec{j}, \text{ where } C : \vec{r}(t) = (t + \sin \frac{\pi t}{2}, t + \cos \frac{\pi t}{2}), 0 \leq t \leq 1.$$

$$(b) \vec{F}(x, y) = \frac{y^2}{1+x^2}\vec{i} + 2y \arctan x\vec{j}, \text{ where } C : \vec{r}(t) = t^2\vec{i} + 2t\vec{j}, 0 \leq t \leq 1.$$

$$(c) \vec{F}(x, y) = (2xz + y^2)\vec{i} + 2xy\vec{j} + (x^2 + 3z^2)\vec{k}, \text{ where } C : x = t^2, y = t + 1, z = 2t - 1, 0 \leq t \leq 1.$$

$$(d) \vec{F}(x, y) = e^y\vec{i} + xe^y\vec{j} + (z + 1)e^z\vec{k}, \text{ where } C : x = t, y = t^2, z = t^3, 0 \leq t \leq 1.$$

## Chapter 6. Surface Integrals

**Exercise 100.** Evaluate the surface integrals of scalar fields.

$$(a) \iint_F xy \, dS, \text{ where } F \text{ is the triangular region with vertices } (1; 0; 0), (0; 2; 0), \text{ and } (0; 0; 2).$$

$$(b) \iint_F yz \, dS, \text{ where } F \text{ is the part of the plane } x + y + z = 1 \text{ that lies in the first octant.}$$

$$(c) \iint_F yz \, dS, \text{ where } F \text{ is the surface with parametric equations } x = u^2, y = u \sin v, z = u \cos v, 0 \leq u \leq 1, \\ 0 \leq v \leq \frac{\pi}{2}.$$

$$(d) \iint_F z \, dS, \text{ where } F \text{ is the surface } x = y + 2z^2, 0 \leq y \leq 1, 0 \leq z \leq 1.$$

$$(e) \iint_F y^2 \, dS, \text{ where } F \text{ is the part of the sphere } x^2 + y^2 + z^2 = 4 \text{ that lies inside the cylinder } x^2 + y^2 = 1 \\ \text{and above the } xy\text{-plane.}$$

$$(f) \iint_F \frac{dS}{(2+x+y+z)^2}, \text{ where } F \text{ is the boundary of the triangular pyramid } x + y + z \leq 1; x \geq 0; y \geq 0; z \geq 0.$$

**Exercise 101.** Find the area of

- (a) the ellipse cut from the plane  $z = 2x + y$  by the cylinder  $x^2 + y^2 = 1$ .
- (b) the surface  $x^2 - 2 \ln x + \sqrt{15}y - z = 0$  above the square  $D = \{(x, y) \in \mathbb{R}^2 : 1 \leq x \leq 2, 0 \leq y \leq 1\}$  in the  $xy$ -plane.
- (c) the part of the paraboloid  $z = x^2 + y^2$  which lies under the plane  $z = 6$ .
- (d) the surface determined by the parametric equations  $x = z(\cos u + u \sin u)$ ,  $y = z(\sin u - u \cos u)$ ,  $0 \leq u, z \leq 1$ .

**Exercise 102.** Find the mass of the surface  $F$  determined by the parametric equations  $x = uv, y = u + v, z = u - v, u^2 + v^2 \leq 1, u \geq 0, v \geq 0$  if the density  $\rho(x, y, z) = x + yz$ .

**Exercise 103.** Find the center of mass of

- (a) a thin hemisphere of radius  $R$  and constant mass density  $C$ .
- (b) the triangle with vertices  $(1; 0; 0)$ ,  $(0; 1; 0)$ ,  $(0; 0; 1)$  and the density  $\rho(x, y, z) = x + 2y + z$ .
- (c) the cylinder  $x^2 + y^2 = 1, 0 \leq z \leq 1$  and the density  $\rho(x, y, z) = x^2 + y^2 + z^2$ .

**Exercise 104.** Evaluate the surface integral  $\iint_A \vec{F} \cdot \vec{n} dS$  for the given vector field  $\vec{F}$  and the oriented surface  $A$ . For closed surfaces, use the positive (outward) orientation.

- (a)  $\vec{F}(x, y, z) = xze^y \vec{i} - xze^y \vec{j} + z \vec{k}$ ,  $A$  is the part of the plane  $x + y + z = 1$  in the first octant and has downward orientation.
- (b)  $\vec{F}(x, y, z) = x \vec{i} + y \vec{j} + z^4 \vec{k}$ ,  $A$  is the part of the cone  $z = \sqrt{x^2 + y^2}$  beneath the plane  $z = 1$  with downward orientation.
- (c)  $\vec{F}(x, y, z) = xz \vec{i} + x \vec{j} + y \vec{k}$ ,  $A$  is the hemisphere  $x^2 + y^2 + z^2 = 25; y \geq 0$ , oriented in the direction of the positive  $y$ -axis.
- (d)  $\vec{F}(x, y, z) = xy \vec{i} + 4x^2 \vec{j} + yz \vec{k}$ ,  $A$  is the surface  $z = xe^y, 0 \leq x, y \leq 1$ , with upward orientation.
- (e)  $\vec{F}(x, y, z) = x^2 \vec{i} + y^2 \vec{j} + z^2 \vec{k}$ ,  $A$  is the boundary of the solid half-cylinder  $0 \leq z \leq \sqrt{1 - y^2}, 0 \leq x \leq 2$ .
- (f)  $\vec{F}(x, y, z) = (x, y, z)$  and  $A$  is the upper surface, upward oriented,  $z = 16 - x^2 - y^2, x^2 + y^2 \leq 16$ .
- (g)  $\vec{F}(x, y, z) = \left(\frac{x}{z}, \frac{y}{z}, z - 2\right)$  and  $A$  is the upper surface, upward oriented, of  $z = 4 - x^2 - y^2, x^2 + y^2 \leq 2$ .
- (h)  $\vec{F}(x, y, z) = (0, y, -z)$  and  $A$  consists of the paraboloid  $y = x^2 + z^2, 0 \leq y \leq 1$  and the disk  $x^2 + z^2 \leq 1, y = 1$ .

**Exercise 105.** Evaluate the surface integral  $\iint_A \vec{F} \cdot \vec{n} dS$  for the given vector field  $\vec{F}$  and the oriented surface  $A$ .

- (a)  $\vec{F}(x, y, z) = (x, z, y)$  and  $A$  is the sphere  $x^2 + y^2 + z^2 = 1$ , oriented outward.

(b)  $\vec{F}(x, y, z) = (x, 2y, 3z)$  and  $A$  is the cube  $[1; 2] \times [1; 2] \times [1; 2]$ , oriented outward.

(c)  $\vec{F}(x, y, z) = (x + 2y, 2y + 3z, 3z + x)$  and  $A$  the triangular pyramid  $ODBC$ ,  $O(0; 0; 0)$ ,  $D(1; 0; 0)$ ,  $B(0; 1; 0)$ ,  $C(0; 0; 1)$ , oriented outward.

**Exercise 106.** A fluid with density  $C$  flows with velocity  $\vec{v} = (yx^2, x, z)$ . Find the rate of flow upward through the paraboloid  $A: z = 9 - \frac{x^2 + y^2}{4}$ ,  $x^2 + y^2 \leq 36$ .

**Exercise 107.** Let  $\vec{F}$  be an inverse square field, that is  $\vec{F}(\vec{r}) = C \frac{\vec{r}}{|\vec{r}|^3}$ , for some constant  $C$ , where  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$ . Show that the surface integral  $\iint_A \vec{F} \cdot \vec{n} dS$ , where  $A$  is a sphere with center at the origin, is independent of the radius of  $A$ .

**Exercise 108.** Use Stokes' Theorem to evaluate  $\int_C \vec{F} \cdot d\vec{r}$ . In each case  $C$  is oriented counterclockwise as viewed from above.

(a)  $\vec{F}(x, y, z) = yz\vec{i} + 2xz\vec{j} + 3xy\vec{k}$  and  $C$  is the circle  $x^2 + y^2 = 4$ ,  $z = 10$ .

(b)  $\vec{F}(x, y, z) = (3x + 2y^2)\vec{i} + (8y + \frac{z^2}{3})\vec{j} + (4z + \frac{3x^2}{2})\vec{k}$  and  $C$  is the boundary of the triangle with vertices  $(2; 0; 0)$ ,  $(0; 3; 0)$  and  $(0; 0; 6)$ .

(c)  $\vec{F}(x, y, z) = xy\vec{i} + 2z\vec{j} + 3y\vec{k}$  and  $C$  is the curve of intersection of the plane  $x + z = 5$  and the cylinder  $x^2 + y^2 = 9$ .

(d)  $\vec{F}(x, y, z) = x^2z\vec{i} + xy^2\vec{j} + z^2\vec{k}$  and  $C$  is the curve of intersection of the plane  $x + y + z = 1$  and the cylinder  $x^2 + y^2 = 9$ .

**Exercise 109.** The work done by the force field  $\vec{F}(x, y, z) = (x^x + z^2, y^y + x^2, z^z + y^2)$  when a particle moves under its influence around the close edge of the part of the sphere  $x^2 + y^2 + z^2 = 4$  that lies in the first octant, in a counterclockwise direction as viewed from above.

**Exercise 110.** Use Stokes' Theorem to evaluate  $\iint_A \text{curl } \vec{F} \cdot \vec{n} dS$ .

(a)  $\vec{F}(x, y, z) = 2y \cos z \vec{i} + e^x \sin z \vec{j} + xe^y \vec{k}$  and  $A$  is the hemisphere  $x^2 + y^2 + z^2 = 9$ ,  $z \geq 0$  oriented upward.

(b)  $\vec{F}(x, y, z) = x^2z^2\vec{i} + y^2z^2\vec{j} + xyz\vec{k}$  and  $A$  is the part of the paraboloid  $z = x^2 + y^2$  that lies inside the cylinder  $x^2 + y^2 = 4$ , oriented upward.

(c)  $\vec{F}(x, y, z) = (xyz, xy, x^2yz)$  and  $A$  consists of the top and the four sides but not the bottom of the cube  $[0; 1] \times [0; 1] \times [0; 1]$ , oriented outward.

(d)  $\vec{F}(x, y, z) = (e^xyz, y^2z, 2z)$ ,  $A$  is the part of the hemisphere  $x^2 + y^2 + z^2 = 9$ ,  $x \geq 0$ , that lies inside the cylinder  $y^2 + z^2 = 4$ , oriented in the direction of the positive  $x$ -axis.

**Exercise 111.** Use the Divergence Theorem to calculate  $\iint_A \vec{F} \cdot \vec{n} dS$ .

- (a)  $\vec{F}(x, y, z) = x^3y\vec{i} - x^2y^2\vec{j} - x^2yz\vec{k}$  and  $A$  is the surface of the solid bounded by the hyperboloid  $x^2 + y^2 - z^2 = 1$  and the planes  $z = -2$  and  $z = 2$ .
- (b)  $\vec{F}(x, y, z) = (\cos z + xy^2)\vec{i} + xe^{-z}\vec{j} + (\sin y + x^2z)\vec{k}$  and  $A$  is the surface of the solid bounded by the paraboloid  $z = x^2 + y^2$  and the plane  $z = 4$ .
- (c)  $\vec{F}(x, y, z) = 4x^3z\vec{i} + 4y^3z\vec{j} + 3z^4\vec{k}$  and  $A$  is the sphere with radius  $R$  and center the origin.
- (d)  $\vec{F}(x, y, z) = z^2x\vec{i} + (y^3 + \sin z)\vec{j} + (x^2z + y^2)\vec{k}$  and  $A$  is the upward oriented top half of the sphere  $x^2 + y^2 + z^2 = 1$ .
- (e)  $\vec{F}(x, y, z) = z^2y^{10}\vec{i} + (4x^2y^3 + \sin z)\vec{j} + (2x^2z + y^2)\vec{k}$  and  $A$  is the outward oriented surface of the cube  $[-1; 1] \times [-1; 1] \times [-1; 1]$ .
- (f)  $\vec{F}(x, y, z) = -xy\vec{i} + 3y^2\vec{j} + 3zy\vec{k}$  and  $A$  is the outward oriented surface of the tetrahedron with vertices  $(0; 0; 0)$ ,  $(1; 0; 0)$ ,  $(0; 1; 0)$ , and  $(0; 0; 1)$ .
- (g)  $\vec{F}(x, y, z) = 6xy^2\vec{i} + 3x^2e^{2z}\vec{j} + 2z^3\vec{k}$  and  $A$  is the outward oriented surface of the solid bounded by the cylinder  $x^2 + y^2 = 4$  and the planes  $z = -1$ , and  $z = 2$ .
- (h)  $\vec{F}(x, y, z) = x^5\vec{i} + \frac{10}{3}x^2y^3\vec{j} + 5zy^4\vec{k}$  and  $A$  is the outward oriented surface of the solid bounded by the paraboloid  $z = 1 - x^2 - y^2$  and the plane  $z = 0$ .